

A ROUTE TO MAGNETIC FIELD REVERSALS: AN EXAMPLE OF AN ABC-FORCED NON-LINEAR DYNAMO

O.M. Podvigina¹

International Institute of Earthquake Prediction Theory
and Mathematical Geophysics,
79 bldg. 2, Warshavskoe ave., 113556 Moscow, Russian Federation

Laboratory of General Aerodynamics, Institute of Mechanics,
Lomonosov Moscow State University,
1, Michurinsky ave., 119899 Moscow, Russian Federation

Observatoire de la Côte d'Azur,
BP 4229, 06304 Nice Cedex 4, France

Submitted to *Geophysical & Astrophysical Fluid Dynamics*
26 January 2003

Abstract

We are investigating numerically the non-linear behaviour of a space-periodic MHD system with ABC forcing. Most computations are performed for magnetic Reynolds numbers increasing from 0 to 60 and a fixed kinematic Reynolds number, small enough for the trivial solution with a zero magnetic field to be stable to velocity perturbations. At the critical magnetic Reynolds number for the onset of instability of the trivial solution the dominant eigenvalue of the kinematic dynamo problem is real. In agreement with the bifurcation theory new steady states with non-vanishing magnetic field appear in this bifurcation. Subsequent bifurcations are investigated. A regime is detected, where chaotic variations of the magnetic field orientation (analogous to magnetic field reversals) are observed in the temporal evolution of the system.

Key words: nonlinear magnetic dynamo, bifurcations, reversals

¹E-mail: olgap@mitp.ru

1 Introduction

In numerical modelling of physical phenomena two approaches are feasible. On the one hand, one can develop a model as detailed, as possible. Then an obvious difficulty is encountered: usually simplifications are necessary, owing to complexity of natural phenomena and/or lack of precise data. Nevertheless, if the model is accurate enough, it displays behaviour similar to that of the underlying physical system, when fixed parameter values equal or close to those in the original system are employed. On the other hand, one can consider a class of approximate models, reflecting only global features of the underlying physical system. Parameter values are varied in order to study how this affects the overall behaviour of the system. The second approach is advantageous in that properties characterising the class of systems are revealed; computationally it is usually less demanding, than the first one.

We follow the second approach to examine a succession of temporal regimes which can occur in magnetohydrodynamic (MHD) systems, when Reynolds numbers are increased. In particular, we are concerned with two questions: Which features of MHD systems are responsible for magnetic field reversals – an attribute of the Earth’s magnetic field? Which sequences of bifurcations can bring an MHD system to such a regime?

A comprehensive review of low-dimensional ODE dynamo models of magnetic field reversals of the Earth, discussed in literature prior to 1994, can be found in Jacobs (1994), Chapter 5. The models were categorised into two distinct classes. In the models of the first class it was assumed that reversals were related to MHD instabilities, triggered by finite-amplitude perturbations of otherwise stable nonlinear dynamos. The perturbations were supposed to be controlled by random processes, and various physical mechanisms of non-MHD nature were proposed as their possible sources. In models of the second class the reversals were a feature of the system, with no external perturbations required to initiate a reversal. The model of solar magnetic field reversals by Zeldovich *et al.* (1983) also falls within this class. Hide *et al.* (1996) and Moroz *et al.* (1998) generalised the Rikitake’s disk model.

Armbruster *et al.* (2001) proposed a 7-dimensional ODE dynamo model obtained by center manifold reduction with the use of symmetry considerations. They considered convection in a spherical shell without rotation for such parameter values, that mode interaction took place and heteroclinic cycles emerged in the phase space, connecting the 7 equilibria present in their system. The heteroclinic cycles were stable due to the symmetries of the system. Introduction of magnetic field could transform the hydrodynamic equilibria into MHD ones, with a non-vanishing magnetic field. The resultant heteroclinic cycles involved changes of orientation of the magnetic field, i.e., reversals of the magnetic dipole (non-dipole magnetic components were ignored in their truncated system).

In the model of Melbourne *et al.* (2001) reversals were also linked with an underlying heteroclinic cycle. To obtain the cycle, they considered con-

vection in a spherical shell with rotation and derived a normal form near a codimension-3 bifurcation, assuming that the flow sustaining the dynamo possessed a certain group of symmetries. The normal form, truncated up to terms of the third order, involved 9 parameters. When several higher-order small symmetry-breaking terms were added in the normal form, for some parameter values the behaviour of the system resembled the Earth's magnetic field reversals.

Reversals were observed in simulations of MHD systems with the geometry and boundary conditions corresponding to those of the Earth in the 3-dimensional model (Glatzmaier and Roberts, 1995, 1996; Roberts and Glatzmaier, 2001), involving the Navier-Stokes equation with the Coriolis, buoyancy and Lorentz forces, the magnetic induction and the temperature equations. A $2\frac{1}{2}$ -dimensional model with a limited resolution in the azimuthal direction, derived from these PDE's, also featured magnetic reversals (Sarson and Jones, 1999; Sarson, 2000). Magnetic field generated in these models was predominantly dipolar; the lengths of time intervals of constant polarity and a short duration of each reversal were consistent with those of the Earth. However, the computations were very demanding in CPU time, prohibiting numerical identification of complex bifurcations in such systems.

In simulations of Glatzmaier and Roberts reversals were robust. Glatzmaier and Roberts (1996) used a modified set of equations (compared to the one studied by Glatzmaier and Roberts, 1995: compositional buoyancy was added and the anelastic approximation was used instead of the Boussinesq approximation), but this did not result in significant changes of the temporal behaviour of magnetic field. Despite the model was run very far from the regime of the geophysical parameter values (Jones, 2000), there was a strong similarity between the model output and the geodynamo. This is another indication that their MHD attractor is structurally stable (i.e., persists when equations, geometry, or parameter values are considerably varied). Roberts and Glatzmaier (2001) considered three different radii of the inner core, corresponding to its present size and some values in the past and in the future (geodynamo was driven by the energy influx due to the latent heat release at the freezing surface of the inner core, slowly growing in time). Again, in all the three cases magnetic field of comparable strength was generated, and the magnetic field exhibited a quantitatively similar temporal behaviour, the system in the future displaying a stronger temporal variability.

2 Statement of the problem

Equations of magnetohydrodynamics are invariant under the symmetry, which preserves flows $\mathbf{v}$ and changes orientation of magnetic fields $\mathbf{b}$:

$$h : (\mathbf{v}, \mathbf{b}) \rightarrow (\mathbf{v}, -\mathbf{b}). \quad (1)$$

One can anticipate the following scenario of development of a temporal regime involving magnetic field reversals. Suppose for certain values of a control parameter the system possesses a stable steady state, in which magnetic field vanishes. When the parameter is varied beyond the critical value, two steady states with a non-vanishing magnetic field emerge in a pitchfork bifurcation (generic to systems with the symmetry (1)). In subsequent bifurcations the steady states bifurcate into more complex attractors, related by (1) and remaining so far distinct. As the parameter is further varied, the attractors of the system join (e.g., due to collision of attractors, or via heteroclinic connection) and a unique attractor emerges. A sample trajectory on the attractor jumps between former attractors (which degenerate into unstable invariant sets). Since the jumps involve changes in the magnetic field orientation, the resulting intermittency can thus be linked with the reversals.

The goal of the present study is to check whether this scenario is feasible. As a test case we consider the Navier-Stokes equation

$$\frac{\partial \mathbf{v}}{\partial t} = \mathbf{v} \times (\nabla \times \mathbf{v}) + \frac{1}{R} \Delta \mathbf{v} - \mathbf{b} \times (\nabla \times \mathbf{b}) + \mathbf{f} - \nabla p \quad (2.a)$$

and the magnetic induction equation

$$\frac{\partial \mathbf{b}}{\partial t} = \nabla \times (\mathbf{v} \times \mathbf{b}) + \frac{1}{R_m} \Delta \mathbf{b} \quad (2.b)$$

under the solenoidality conditions

$$\nabla \cdot \mathbf{v} = 0, \quad \nabla \cdot \mathbf{b} = 0. \quad (2.c)$$

Here $\mathbf{v}$ is velocity of the flow, $\mathbf{b}$ – magnetic field, p – pressure, R and R_m are kinematic and magnetic Reynolds numbers. (Since in the problem under consideration characteristic length and velocity magnitudes are of order one, the Reynolds numbers are defined as inverse viscosity and inverse magnetic diffusivity, respectively.) The fields are supposed to be 2π -periodic in space.

The force

$$\mathbf{f} = \mathbf{u}_0 / R \quad (3)$$

is assumed, where $\mathbf{u}_0$ is an ABC flow

$$\mathbf{u}_0 = (A \sin x_3 + C \cos x_2, B \sin x_1 + A \cos x_3, C \sin x_2 + B \cos x_1). \quad (4)$$

For this force,

$$\mathbf{v} = \mathbf{u}_0, \quad \mathbf{b} = 0 \quad (5)$$

is a steady solution to (2)-(4) for all Reynolds numbers.

This particular system is considered for the following reasons: First, space periodicity enables one to use pseudo-spectral methods (see Canuto *et al.*, 1989; Boyd, 1989), which are computationally efficient. Second, linear stability of the system (2)-(5) to hydrodynamic (Galloway and Frisch, 1987; Podvigina

and Pouquet, 1994; Podvigina, 1999) and magnetic (Arnold and Korkina, 1983; Galloway and Frisch, 1986; Galanti *et al.*, 1992; Childress and Gilbert, 1995) perturbations has been investigated both numerically and analytically (nonlinear regimes for some parameter values were also explored by Podvigina and Pouquet, 1994; Podvigina, 1999; Galanti *et al.*, 1992; Feudel *et al.*, 1996; Brummell *et al.*, 2001). These studies provide guidance, in what parameter ranges the targeted type of behaviour may be observed. In particular, growing magnetic modes exist for the flow (3) in broad intervals of constants A , B and C . Third, since the critical magnetic Reynolds number is typically of the order of 10, only moderate R_m need to be considered, and a relatively low resolution (32^3 Fourier harmonics) suffices.

Magnetic field growth rates were calculated by Galanti *et al.* (1992) for the flow (4) with the coefficients satisfying $B = C$, $A^2 + B^2 + C^2 = 3$, $0 < B/A < 1$ for various values of R_m . For $R_m = 12$ there are three windows in B/A of positive growth rates (see Fig. 2 *ibid.*). We have checked that in the window $0.7 < B/A < 0.9$ the dominant eigenvalue is real, and in the other two windows it is complex. Distinct steady states with a non-zero magnetic field appear in a bifurcation of the trivial steady state (5), only if the magnetic induction operator has a zero (hence real) eigenvalue at the bifurcation point. This has suggested to set

$$A = 1, \quad B = C = 0.75; \quad (6)$$

these values of constants in (4) are assumed throughout.

With our goal in mind, in the present study we focus at non-hydrodynamic attractors. The hydrodynamic global stability of $\mathbf{u}_0$ guarantees that magnetic field does not vanish in saturated regime, if $\mathbf{v} = \mathbf{u}_0$ is a kinematic dynamo. We have verified that for $R = 4$, $\mathbf{v} = \mathbf{u}_0$ is a unique attractor of the hydrodynamic system ((2.a,c) with $\mathbf{b} = 0$), and most computations are made for this value of R . Several runs are also performed for higher R .

3 The group of symmetries

The group of symmetries without inversion of time of an ABC flow with $B = C$, which we denote by $\mathcal{G}$, is comprised of two independent symmetries

$$s_1 : x_1 \rightarrow -\frac{\pi}{2} - x_3, \quad x_2 \rightarrow -\frac{\pi}{2} - x_2, \quad x_3 \rightarrow -\frac{\pi}{2} - x_1,$$

$$s_2 : x_1 \rightarrow \frac{\pi}{2} + x_3, \quad x_2 \rightarrow -\frac{\pi}{2} + x_2, \quad x_3 \rightarrow \frac{\pi}{2} - x_1,$$

their superpositions

$$s_3 = s_2^2 : x_1 \rightarrow \pi - x_1, \quad x_2 \rightarrow \pi + x_2, \quad x_3 \rightarrow -x_3,$$

$$s_4 = s_2^3 : x_1 \rightarrow \frac{\pi}{2} - x_3, \quad x_2 \rightarrow \frac{\pi}{2} + x_2, \quad x_3 \rightarrow -\frac{\pi}{2} + x_1,$$

$$s_5 = s_1 s_2 : x_1 \rightarrow \pi + x_1, \quad x_2 \rightarrow -x_2, \quad x_3 \rightarrow \pi - x_3,$$

$$s_6 = s_1 s_2^2 : x_1 \rightarrow -\frac{\pi}{2} + x_3, x_2 \rightarrow \frac{\pi}{2} - x_2, x_3 \rightarrow \frac{\pi}{2} + x_1$$

$$s_7 = s_1 s_2^3 : x_1 \rightarrow -x_1, x_2 \rightarrow \pi - x_2, x_3 \rightarrow \pi + x_3,$$

and the identity transformation $s_8 = e$ (Arnold, 1984; Dombre *et al.*, 1986; Podvigina and Pouquet, 1994). The group is isomorphic to the symmetry group of a square, $\mathbf{D}_4$.

Any symmetry of the force (3), (4) which does not involve inversion of time is also a symmetry of the Navier-Stokes equation (2.a) with $\mathbf{b} = 0$. The group of symmetries of the MHD system (2)-(4), (6), which we denote by $\mathcal{H}$, is a direct product of $\mathcal{G}$ and the 2-element group $\mathbf{Z}_2$, generated by h (1). It has 16 elements, which are either s_i , or hs_i .

Problems in hydrodynamics and magnetohydrodynamics (see e.g. the review by Crawford and Knobloch, 1991) often involve symmetries, induced by the geometry of the region where fluid is contained. In particular, if convection in a plane layer is restricted to space-periodic flows with equal periods along two Cartesian axes parallel to boundaries, and conditions at the boundaries are different, the group of symmetries of the system is isomorphic to $\mathcal{G}$. If in addition magnetic field is present, the symmetry group is isomorphic to $\mathcal{H}$. Thus, though the system under consideration (2)-(4), (6) is highly symmetric, it can not be regarded as an exotic one.

Symmetries are useful for categorisation of attractors. Let $\mathcal{A}$ be an attractor of a dynamical system, invariant under a symmetry g : $g(\mathcal{A}) = \mathcal{A}$. Two cases can be distinguished: either $\mathcal{A}$ is pointwise invariant, i.e. $g(\mathbf{x}) = \mathbf{x}$ for all points $\mathbf{x} \in \mathcal{A}$, or it is invariant only as a set, with $g(\mathbf{x}) \neq \mathbf{x}$ for some $\mathbf{x} \in \mathcal{A}$. In what follows, only the symmetries for which an attractor is pointwise invariant are regarded as symmetries of the attractor.

4 Attractors of the MHD system for $R = 4$

Results of computations are summarised in Table 1 (see also the bifurcation diagram Fig.1). The magnetic Reynolds number has been increased step 1, except in the interval $37 < R_m < 40$, where a high density of bifurcations has required smaller steps. For every considered R_m an attractor has been obtained in a run with an initial condition, which is a perturbation of the trivial steady state (5) with the energy of 10^{-6} in each Fourier harmonics spherical shell. The range of existence of attractors has been determined by continuation in parameter: the runs are done with an initial condition, which is a point on the attractor for close values of R_m .

For $15 \leq R_m \leq 58$ the system under consideration possesses multiple attractors. For $16 \leq R_m \leq 58$ attractors, which we have detected, belong to two independent families; within each family they are genetically related by sequences of bifurcations. Attractors from different families have distinct groups of symmetries. The families can be distinguished also by their time-averaged

Table 1. Attractors, detected for the MHD system with the forcing (3), (4), (6) for $R = 4$ and $0 \leq R_m \leq 60$. The third column shows the number of elements of the symmetry group for which an attractor is pointwise invariant, and the fourth column – generators of the group. Indices in the labels of attractors (second column) have the following meaning. For periodic orbits emerging in the period doubling cascade the superscript M refers to an $M\tau$ -period orbit, τ being the period of the foremost orbit. The first subscript labels different attractors of a given kind. The second subscript labels mutually symmetric attractors.

R	Attractors	Number of symmetries	Generators	$\overline{E}_k$	$\overline{E}_m$
$R_m \leq 14$	steady state S_0	16	s_1, s_2, h	1.06	0
$15 \leq R_m \leq 24$	steady states $S_{1,i}$ ($i = 1, 2$)	8	hs_1, s_2	1.06-0.92	0-0.035
$25 \leq R_m \leq 37$	periodic orbits $P_{1,i}$ ($i = 1, 2$)	4	$hs_1, s_3 (= s_2^2)$	0.94	0.03
$37.5 \leq R_m \leq 38.5$	steady states $S_{1,i}$ ($i = 1, 2$)	8	hs_1, s_2	0.95	0.025
$39 \leq R_m \leq 39.2$	periodic orbits $P_{2,i}^1$ ($i = 1, 2$)	8	hs_1, s_2	0.98	0.02
$39.3 \leq R_m \leq 39.4$	periodic orbits $P_{2,i}^2$ ($i = 1, 2$)	8	hs_1, s_2	0.98	0.02
$R_m = 39.45$	periodic orbits $P_{2,i}^4$ ($i = 1, 2$)	8	hs_1, s_2	0.98	0.02
$R_m = 39.5$	chaotic $C_{1,i} = P_{2,i}^\infty$ ($i = 1, 2$)	8	hs_1, s_2	0.98	0.02
$R_m = 39.6$	periodic orbits $P_{2,i}^1$ ($i = 1, 2$)	8	hs_1, s_2	0.97	0.025
	& periodic orbit P_3	8	hs_1, s_2	0.95	0.03
$39.7 \leq R_m \leq 41$	periodic orbit P_4	8	hs_1, s_2	0.93	0.035
$R_m = 42, 43, 44$	tori $T_{1,i}$ ($i = 1, 2$)	4	hs_1, s_3	0.93	0.04
$R_m = 45$	chaotic $C_{2,i}$ ($i = 1, 2$)	4	hs_1, s_3	0.93	0.04
$R_m = 46, 47$	chaotic $C_{3,i}$ ($i = 1, 4$)	2	hs_1	0.94	0.03
$R_m = 48, 49$	chaotic $C_{4,i}$ ($i = 1, 4$)	2	hs_1	0.92	0.035
$R_m = 50, 51$	chaotic $C_{5,i}$ ($i = 1, 2$)	2	hs_1	0.91	0.04
$52 \leq R_m \leq 58$	tori $T_{2,i}$ ($i = 1, 4$)	2	hs_1	0.9	0.05
$16 \leq R_m \leq 25$	periodic orbit P_6	4	hs_2	0.4-0.7	0.07-0.2
$26 \leq R_m \leq 60$	torus T_3	4	hs_2	0.3-0.4	0.2-0.25

kinetic ($\overline{E}_k$) and magnetic ($\overline{E}_m$) energies: $0.9 < \overline{E}_k < 1.1$, $0 \leq \overline{E}_m < 0.05$ for the first family, and $0.3 < \overline{E}_k < 0.7$, $0.07 < \overline{E}_m < 0.25$ for the second one.

The first family exists for $0 \leq R_m \leq 58$: the first attractor is the trivial steady state (5) (denoted by S_0) remaining stable up to² $R_m = 14$. It becomes unstable in a pitchfork bifurcation, in which two stable mutually symmetric steady states ($S_{1,1}$ and $S_{1,2}$) with a non-zero magnetic field emerge. The 8-element symmetry group of $S_{1,i}$ is isomorphic to $\mathbf{D}_4$; however, it is distinct from $\mathcal{G}$.

The steady states $S_{1,i}$ become unstable in a supercritical Hopf bifurcation at the interval $24 < R_m < 25$. A stable periodic orbit $P_{1,i}$ of period ≈ 12 appears in a vicinity of the steady state $S_{1,i}$. The two orbits are mutually symmetric (they are interrelated by h). Each periodic orbit possesses all 8 symmetries of its parent steady state, but each individual point of the orbit has a symmetry group of only 4 elements (it is isomorphic to $\mathbf{D}_2$). The orbits exist for $25 \leq R_m \leq 37$ remaining attracting. A typical behaviour of a trajectory is shown on Fig. 2 for $R_m = 30$. After an initially small magnetic energy at first exponentially grows and afterwards decays with oscillations, it subsequently levels off (see a plateau about 400 time units long on Fig. 2a): the trajectory in the phase space evolves in a vicinity of $S_{1,i}$ (which is unstable now). In the further evolution the trajectory leaves the steady state, being attracted by $P_{1,i}$ (see Fig. 2b). The orbits disappear in a subcritical Hopf bifurcation between $R_m = 37$ and $R_m = 37.5$. For $R_m = 37.5$, 38 and 38.5 the two steady states $S_{1,i}$ are verified to be stable.

Next bifurcation of $S_{1,i}$ is again a supercritical Hopf bifurcation. The emerging mutually symmetric periodic orbits $P_{2,i}^1$, $i = 1, 2$ (one for each of the two steady states) possess all the symmetries of the steady states. For $39 \leq R_m \leq 39.2$ they are attracting, and their period is $\tau(P_{2,i}^1) \approx 120$. Behavior of a sample trajectory in the phase space (see Fig. 3) resembles the one shown on Fig. 2; however, on Fig. 3a the plateau corresponding to the evolution in the vicinity of the unstable steady state $S_{1,i}$ virtually disappears, and the period is much larger. The orbit is located in a different region of the phase space (cf. Fig. 2b and Fig. 3b). At an R_m between 39.2 and 39.3 two mutually symmetric orbits $P_{2,i}^2$ ($i = 1, 2$) of a twice larger period emerge (see Fig. 4a). A sequence of period-doubling bifurcations begins. The next period doubling occurs at the interval $39.4 < R_m < 39.45$ (see period-four orbit $P_{2,i}^4$ for $R_m = 39.45$ on Fig. 4b). At $R_m = 39.5$ a trajectory, initially close to S_0 , is either already chaotic, or close to an orbit of a very long period $P_{2,i}^\infty$ (see Fig. 4c). This indicates that at $R_m = 39.5$ the period-doubling cascade is over. For $R_m = 39.6$ a trajectory with the same initial condition is attracted to a period-one orbit $P_{2,i}^1$.

Another attractor exists for $R_m = 39.6$. It reveals itself when the attractor

²This is consistent with results of Galanti *et al.* (1992): since the ABC coefficients (6) are not normalised to satisfy $A^2 + B^2 + C^2 = 3$ as they were in the cited paper, Reynolds numbers referred to here are lower by a factor $(17/24)^{1/2}$.

for $R_m = 39.7$, which is unique in this region of the phase space (see below) is continued in smaller R_m (no attractor of this type is observed by further continuation in smaller R_m). This is a periodic orbit P_3 , involving transitions between the two period-two orbits $P_{2,1}^2$ and $P_{2,2}^2$ (see Fig. 5a). The orbit is invariant under h and any symmetry from $\mathcal{G}$. In the course of temporal evolution, direction of magnetic field is reversed, behaviour of non-zero magnetic field Fourier coefficients is similar to that shown on Fig. 5b. A reversal takes a relatively short time, during which initially the energy of magnetic field attains the minimum and kinetic – the maximum, and subsequently magnetic energy blows up to a maximum, with the kinetic energy simultaneously reaching its minimum (Fig. 5c,d). Thereby, during a reversal magnetic energy is at first transformed into kinetic one, and then back into magnetic energy; for higher Reynolds numbers reversals exhibit such energy transformations as well.

This bifurcation is apparently of the following nature. In the period-doubling cascade infinitely many unstable periodic orbits $P_{2,i}^M$ ($i = 1, 2$) of periods $\tau(P_{2,i}^M) = M\tau(P_{2,i}^1)$ are created. Consider two such orbits $P_{2,i}^m$ ($m = 2$ in our case) related by h . Suppose that for some value of a control parameter the stable manifold of $P_{2,1}^m$ intersects the unstable manifold of $P_{2,2}^m$, and a heteroclinic trajectory connecting these orbits emerges. Consider the symmetry, which maps one of the periodic orbits to another. Under this symmetry the heteroclinic trajectory is mapped to a distinct heteroclinic trajectory, and the two form a heteroclinic cycle. The cycle may give rise to a periodic orbit P_3 , when the control parameter varies. (This scenario is analogous to that in two-dimensional dynamical systems, where a homoclinic connection generically implies for close values of the control parameter existence of periodic orbits in a vicinity of the homoclinic trajectory, and their period tends to infinity when the critical value is approached.) Thus, the regime of reversals emerges via heteroclinic connection of two periodic orbits.

For $39.7 \leq R_m \leq 41$ sample trajectories, initially close to the trivial steady state S_0 , are attracted by a periodic orbit P_4 . While P_3 (for $R_m = 39.6$) follows closely each of $P_{2,i}^2$ (Fig. 5a), this is not the case any more for P_4 (for the higher R_m 's; see Fig. 6). In agreement with our conjecture that emergence of the orbit is related to heteroclinic connection, the period decreases from 800 for $R_m = 39.7$ to 500 for $R_m = 41$.

For $39.6 \leq R_m \leq 41$ the attractor involving reversals is unique. Behaviour of all magnetic field Fourier harmonics is coherent and resembles the one shown on Fig. 5b: time intervals, where a Fourier coefficient remains close to a certain value, are separated by jumps changing the sign of the coefficient, but not its magnitude. In contrast with the coherent behaviour of all magnetic field Fourier harmonics for $39.6 \leq R_m \leq 41$, they can be divided for $42 \leq R_m \leq 51$ into two groups with a qualitatively different behaviour. Behaviour of harmonics from the first group is the same as above, and hence the system exhibits reversals. Harmonics from the second group oscillate with a relatively short temporal period, some of their time averages do not vanish. The total time-averaged energy of harmonics from the first group is higher than that from

the second group (for instance, cf. Fig. 7a and 7b). The Earth's magnetic field shows a similar behavior: Spherical harmonics can be categorized into the dipole and the quadrupole families according to their symmetry about the equatorial plane (Roberts and Stix, 1972), and paleomagnetic data suggest that a magnetic reversal in the Earth is more likely to occur, when the ratio of the magnitude of the quadrupole-family component of the magnetic field to that of the dipole-family component is high (Merrill and McFadden, 1988). For $42 \leq R_m \leq 51$ our attractors have no the symmetry h on average (since the time average of some harmonics is non-zero), and thus h maps an attractor to a different attractor with reversals. Again, a parallel can be drawn with a property of the magnetic field of the Earth: Paleomagnetic data indicate that the time-averaged magnetic field of the Earth lacks the north-south symmetry and there are significant (though small) differences between the states of the normal and reverse polarity (Merrill *et al.*, 1979), this implying the absence of the symmetry h on average.

For $R_m = 42$ and 43 the regime is quasi-periodic: the orbit P_4 bifurcates into two attracting tori $T_{1,i}$ ($i = 1, 2$) interrelated by h (see Fig. 7). The second frequency can be observed on Fig. 7a, showing real part of the Fourier component $b_{0,1,1}^3$ of magnetic field. The time average of this Fourier component is negative (Fig. 7a); hence, the symmetry h maps this attractor to a distinct one. (For $37.5 \leq R_m \leq 41$ the component vanishes, because former attractors [P_4 for $39.7 \leq R_m \leq 41$, P_3 for $R_m = 39.6$, $P_{2,i}^m$ for $39 \leq R_m \leq 39.6$ and $S_{1,i}$ for $37.5 \leq R_m \leq 38.5$] possess the symmetry $hs_5 = hs_1s_2$.)

For $R_m = 45$ two mutually symmetric attractors persist (Fig. 8): each torus $T_{1,i}$ bifurcates into a chaotic attractor $C_{2,i}$. Reversals are less regular, yet they are too ordered in comparison with those of the Earth's magnetic field.

For $R_m = 46$ (Fig. 9) behaviour of a sample trajectory suggests existence of four (unstable) steady states $S_{2,i}$ (see plateaux at $1500 \leq t \leq 2100$). They are mutually symmetric, and have the same 4-element symmetry group generated by hs_1 and s_3 . Apparently two $S_{2,i}$'s emerged from each $S_{1,i}$ in a pitchfork bifurcation. Initially (at $0 \leq t < 700$) the sample trajectory for $R_m = 46$ undergoes several reversals, similar in nature to those observed for $39.7 \leq R_m \leq 45$. Afterwards it is attracted to a steady state $S_{2,i}$. The behaviour near the steady state resembles the Shilnikov attractor: the trajectory approaches the steady state along the two-dimensional eigenspace associated with the complex eigenvalues with the maximal negative real part, and leaves it along the one-dimensional unstable manifold. Subsequent sample evolution consists of transitions between the steady state $S_{2,i}$ and the region of the phase space where a former chaotic attractor $C_{2,i}$ was located (duration of repeating events in the saturated regime is ≈ 1700). In particular, large-amplitude excursions at $2350 \leq t \leq 2500$, $2300 \leq t \leq 2450$, $4000 \leq t \leq 4150$, etc., are reminiscent of the behaviour of a $C_{2,i}$ trajectory for $R_m = 45$. Thus, apparently two former chaotic attractors $C_{2,i}$ have disappeared in collision with the four steady states $S_{2,i}$ to give rise to four new chaotic attractors $C_{3,i}$. It is

notable that the sample trajectory leaves a vicinity of the $S_{2,i}$ in alternating directions along the unstable manifold. For $R_m = 47$ the behaviour is similar to the one at $R_m = 46$.

For $R_m = 48$ the system possesses four mutually symmetric attractors $C_{4,i}$. Events are longer than in the previous regime, about 3000 time units (see Fig. 10). They exhibit a new feature, a phase of initially exponentially decaying oscillations (e.g. at $1700 \leq t \leq 2300$ and $4500 \leq t \leq 5200$). This indicates existence of a weakly unstable periodic orbit of a new kind, $P_{5,i}$ ($i = 1, 4$), in a vicinity of the former chaotic attractor $C_{3,i}$. A similar behaviour is also observed for $R_m = 49$.

For $R_m = 50, 51$ (Fig. 11) the four $C_{4,i}$ are superceded by two new attractors, $C_{5,i}$. Large amplitude oscillations, e.g. at $7500 \leq t \leq 8500$, are due to attraction of the trajectory by weakly unstable tori $T_{2,i}$ ($i = 1, 4$), which have bifurcated from the periodic orbits $P_{5,i}$. In the saturated regime behaviour consists of three phases: (i) a trajectory is close to a periodic orbit $P_{5,i}$ (e.g. at $2300 \leq t \leq 3000$ and $4300 \leq t \leq 5000$); (ii) the trajectory evolves in the vicinity of a torus $T_{2,i}$ (e.g. at $3200 \leq t \leq 4000$ and $5300 \leq t \leq 6000$); (iii) the trajectory abruptly jumps toward the second periodic orbit $P_{5,i'}$ to reproduce the sequence of phases.

For $R_m = 52$ each of the two former chaotic attractors splits into two attracting tori $T_{2,i}$ (Fig. 12; cf. Figs. 11b and 12b). The four new attractors are mutually symmetric, they are stable for $52 \leq R_m \leq 58$.

For $57 \leq R_m \leq 60$ a sample trajectory initially close to S_0 is attracted by the torus T_3 , belonging to the second family. The family emerges between $R_m = 15$ and $R_m = 16$: in addition to the steady states $S_{1,i}$, for $R_m \geq 16$ the system possesses another attractor, a stable periodic orbit P_6 . The orbit does not exist for $R_m \leq 15$; apparently it appears in a saddlenode bifurcation. For $R_m = 26$ the orbit is unstable and a torus T_3 has appeared in a Hopf bifurcation. T_3 remains an attractor for all higher considered R_m (specifically, for $R_m = 26, 27, 30, 40, 57$ and 60). For $R_m \geq 57$ the torus attracts trajectories (in the phase space) initially close to the trivial steady state S_0 , and for $R_m \geq 59$ it is the only global attractor of the system. The torus T_3 and the orbit P_6 were traced back from $R_m = 57$ by continuation in R_m .

Studying numerically a dynamical system, one can never guarantee that all attractors are found. The computations are intensive (a run with the resolution 32^3 Fourier harmonics up to $t = 5000$ has required several dozens hours of one processor), prohibiting to carry out more detailed computations. For instance, if attractors of two or more new types emerge for a particular considered R_m , in principle we could have missed some of them; however, it is unlikely that several attractors of different morphology are born at the same or close values of the magnetic Reynolds number. In our estimations all attractors, whose basin of attraction contains the trivial steady state, have been identified – up to a natural limitation: since a period-doubling cascade involves an infinite number of bifurcations, it is impossible to capture numerically *all* bifurcations within the cascade. Attractors, which do not attract small per-

turbations of the trivial steady state and which are far from it in the phase space, (if such attractors exist in our system) are not involved in the scenario of emergence of magnetic field reversals, and they are out of the scope of the present investigation.

5 Attractors of the MHD system for $R_m = 40$

Simulations are performed for $R_m = 40$ and for several values of R with an initial condition, same for all runs, being a small perturbation of S_0 (see Table 2). The question we address is how the temporal behaviour changes with R , in particular, if for higher R reversals persist, or whether magnetic field vanishes in a saturated regime. We do not attempt to identify all bifurcations occurring when R varies. Computations with the same R and the same initial condition for the flow are also performed for the purely hydrodynamic system (2.a), in order to compare the behaviour and attractors of the two systems. (If (2) possesses attractors with $\mathbf{b} = 0$ different from the trivial steady state, an initially growing magnetic field can die out in the nonlinear regime, the resulting attractor being purely hydrodynamical, see Brummell *et al.*, 1998; Matthews, 2000.)

For $R = 3$ the detected attractor is the torus T_1 (which is an attractor for $R = 4$ and $26 \leq R_m \leq 60$). For $R = 6$ reversals take place, similar to those observed for $R = 4$ and $R_m = 45$, and the attractor is $C_{2,i}$. Thus, for R close to 4 no new attractors are found.

For $R = 10$ a new attractor was found in the full MHD system – a periodic orbit P_7 with a symmetry group of 8 elements. Comparison of Tables 2 and 3 reveals no relation of attractors of the hydrodynamic and MHD systems for $R = 10, 15$ and 20. For these Reynolds numbers a small magnetic field (magnetic energy E_m is below 0.07; see Table 2) drastically changes behaviour of the system – the hydrodynamic and MHD systems have attractors of different types, and the average kinetic energy $\overline{E}_k$ decreases significantly, e.g. $\overline{E}_m = 0.01$ and $\overline{E}_k$ decreases from 0.9 (the hydrodynamic case) to 0.7 (the MHD case) for $R = 20$.

For $R = 15$ behaviour of the sample trajectory of the MHD system in saturated regime is chaotic. But before the saturated regime sets in, the trajectory is attracted by P_7 and for $1000 \leq t \leq 2000$ remains close to this periodic orbit (see Fig. 13), which is now weakly unstable. For $R = 20$ the trajectory with the same initial condition is attracted to a new torus $T_{4,i}$; initially the temporal behaviour of the trajectory resembles the one observed for $R = 15$ (cf. Figs. 14 and 14).

For $R = 25$ the MHD system possesses new chaotic attractors $C_{7,i}$ ($i = 1, 2$) with a trivial symmetry group. They resemble the chaotic attractor C_6 observed for $R = 15$ (cf. Fig. 13 and Fig. 15), but unlike $C_{7,i}$, C_6 is unique and has a symmetry group of four elements. Comparison of $C_{7,i}$ with the attractor

Table 2. Attractors, detected for the MHD system with the force (3), (4), (6) for $R_m = 40$ and $3 \leq R \leq 25$. The third column shows the number of elements of the symmetry group for which an attractor is pointwise invariant, and the fourth – generators of the group.

R	Attractors	Number of symmetries	Generators	$\overline{E}_k$	$\overline{E}_m$
$R = 3$	T_3	4	hs_2	0.4	0.25
$R = 4$	periodic orbit P_4	8	hs_1, s_2	0.95	0.03
$R = 6$	chaotic $C_{2,i}$ ($i = 1, 2$)	4	hs_1, s_3	0.95	0.03
$R = 10$	periodic orbit P_7	8	hs_1, s_2	0.5	0.06
$R = 15$	chaotic C_6	4	s_1, s_3	0.6	0.04
$R = 20$	tori $T_{4,i}$ ($i = 1, 2$)	4	s_1, s_3	0.7	0.01
$R = 25$	chaotic $C_{7,i}$ ($i = 1, 2$)	1	e	0.5	0.02

Table 3. Attractors, detected for the hydrodynamic system (2.a) with the force (3), (4), (6) for $3 \leq R \leq 25$. The third column shows the number of elements of the symmetry group for which an attractor is pointwise invariant, and the fourth column – generators of the group.

R	Attractors	Number of symmetries	Generators	$\overline{E}_k$
$R = 3, 4, 6$	$\mathbf{u}_{ABC}$	8	s_1, s_2	1.06
$R = 10^*$	$\mathbf{u}_{ABC}$	8	s_1, s_2	1.06
	S_1^h	8	s_1, s_2	0.99
$R = 15$	S_1^h	8	s_1, s_2	0.93
$R = 20$	chaotic C_1^h	4	s_2	0.9
$R = 25$	chaotic C_2^h	1	e	0.7

* S_1^h for $R = 10$ was traced back from $R = 15$; it is not observed for $R = 6$.

of the non-magnetic Navier-Stokes equation, which also is chaotic, reveals that the influence of magnetic field is in some sense stabilising: Fourier coefficients of the flow experience fewer jumps and the amplitude of their oscillations is much smaller (compare Fourier components of flows on Figs. 15b and 16b). An exponential growth of the initially small magnetic field in the sample evolution of the system begins only at $t = 200$ (Fig. 15c). Accordingly, until $t = 300$ the flow evolution is similar to the one in the absence of magnetic field. Magnetic field starts growing, when the trajectory is attracted by a weakly unstable periodic orbit in the hydrodynamic subspace. Departure from this orbit causes an initial decay of magnetic field before the onset of saturated behaviour.

For $R = 15$ and 20 behaviour of Fourier components of magnetic field is similar to the one shown on Fig. 15d. It consists of chaotic irregular small-period oscillations about zero, different from the behaviour observed for $R = 4$. Thus for $R_m = 40$, $15 \leq R \leq 25$ the considered system is not in a regime resembling reversals.

6 Conclusion

The bifurcation scenario leading to emergence of reversals, which we put forward, proves feasible. Reversals are found for small R , for which the hydrodynamic system has a unique globally stable steady state. Complexity of the sequence of bifurcations obtained in simulations is comparable to that of the hydrodynamic system examined by Podvigina and Pouquet (1994) and Podvigina (1999). Numerous types of behaviour are identified, which involve a large variety of attractors and unstable invariant sets of various types affecting behaviour of evolutionary solutions. The regime of reversals arises as a result of merging of two distinct attractors, apparently via heteroclinic connection due to intersection of stable and unstable manifolds of periodic orbits (in contrast to the system of Armbruster *et al.* (2001), where a heteroclinic cycle emerged simultaneously with magnetic steady states).

Other regimes with a non-zero magnetic field are observed when any of the Reynolds numbers is increased. Thus in our computations reversals are not so robust, as in simulations of Glatzmaier and Roberts. The range of parameters considered here is rich in bifurcations: for $R = 4$ we found 14 bifurcations in the interval $14 < R_m < 60$ (counting the infinite number of period-doubling bifurcations as one), 7 distinct types of attractors involve magnetic field reversals. In another dissimilarity with the results of Glatzmaier and Roberts (and with the Earth's magnetic field) in our simulations $\overline{E}_m < \overline{E}_k$ (cf. $\overline{E}_m/\overline{E}_k \sim 10^4$ for the geodynamo). In our computations reversals appear more regular than those in the Earth, since much smaller values of Reynolds numbers are considered. The dissimilarity with reversals in the Earth also owes to the fact that we solve a simplified system of PDE's (neither an account of thermal or sedimentation-driven convection is taken, nor that of the Coriolis force) in an idealised space-periodic geometry. Nevertheless, it appears possible to mim-

ick the presence of two time scales (average periods of constant polarity are much longer than the average time of a reversal itself); instances of excursions of magnetic field (i.e., incomplete reversals) can be observed as well (see e.g. Figs. 8a and 10).

Our results show that magnetic field reversals are an inherent feature of a nonlinear dynamical system of the MHD type. The fundamental reason for their existence is the symmetry h , changing polarity of the magnetic field. Neither external triggering random processes, such as the ones postulated in earlier studies, nor symmetry breaking due to inhomogeneity of the outer core boundaries, nor rotation, nor a specific heteroclinic structure or symmetries of the flow sustaining a non-linear dynamo are required to give rise to a sequence of reversals. Such physical processes, as heat transfer or sedimentation are also unnecessary *per se* (as soon as the system is under the action of any body force providing input of energy).

Acknowledgments

I am grateful to V. Zheligovsky for discussions. Part of this research was carried out during my stays at the School of Mathematical Sciences, University of Exeter, UK, in January – February 2001 and in May – July 2002. I am grateful to the Royal Society for their support of the two visits. Some numerical results were obtained using computational facilities provided by the program “Simulations Interactives et Visualisation en Astronomie et Mécanique (SIVAM)” at Observatoire de la Côte d’Azur, France. My research work at Observatoire de la Côte d’Azur was supported by the French Ministry of Education. Part of the work was done during a visit to Departamento de Matemática Aplicada, Faculdade de Ciências, University of Porto, Portugal at the kind invitation of Centro de Matemática Aplicada in June – July, 2001. Last but not least, I am grateful to referees for stimulating remarks.

References

- Armbruster D., Chossat P., and Oprea I. Structurally stable heteroclinic cycles and the dynamo dynamics, In: *Dynamo and Dynamics, a Mathematical Challenge* (Eds. Chossat, P., Armbruster, D. and Oprea, I.), 313–322, Kluwer Academic Publishers (2001).
- Arnold V.I. and Korkina E.I. The growth of a magnetic field in a three-dimensional incompressible flow, *Vestnik Moscow State Univ., Ser. Math.* **3**, 43–46 (1983) (in Russian).
- Arnold V.I. On the evolution of magnetic field under the action of advection and diffusion, in *Some problems of modern analysis* (ed. V.M. Tikhomirov). Moscow Univ. Press, 8–21 (1984) (in Russian).
- Boyd J.P. *Chebyshev & Fourier Spectral Methods*. Springer-Verlag, Berlin (1989).
- Brummell N.H., Cattaneo F. and Tobias S.M. Linear and nonlinear dynamo action, *Phys. Lett. A* **28**, 437–442 (1998).
- Brummell N.H., Cattaneo F. and Tobias S.M. Linear and nonlinear dynamo properties of time-dependent ABC flows, *Fluid Dyn. Res.* **28**, 237–265 (2001).
- Canuto C., Hussaini M.Y., Quarteroni A. and Zang T.A. *Spectral Methods in Fluid Dynamics*, Springer-Verlag, Berlin (1988).
- Childress S. and Gilbert A.D. *Stretch, twist, fold: the fast dynamo*. Springer-Verlag, Berlin (1995).
- Crawford J.D. and Knobloch E. Symmetry and symmetry-breaking bifurcations in fluid dynamics, *Annu. Rev. Fluid Mech. A* **23**, 341–387 (1991).
- Dombre T., Frisch U., Greene J.M., Hénon M., Mehr A. and Soward A. Chaotic streamlines in the ABC flows, *J. Fluid Mech.* **167**, 353–391 (1986).
- Feudel S., Seehafer N., Galanti B. and Schmidtman O. Symmetry breaking bifurcation for the magnetohydrodynamic equation with helical forcing, *Phys. Rev. E* **54**, 2589–2596 (1996).
- Galanti B., Sulem P.L. and Pouquet A. Linear and Non-Linear Dynamos Associated With ABC Flows, *Geophys. Astrophys. Fluid Dynamics* **66**, 183–208 (1992).
- Galloway D.J. and Frisch U. Dynamo action in a family of flows with chaotic streamlines, *Geophys. Astrophys. Fluid Dynamics* **36**, 53–83 (1986).
- Galloway D.J. and Frisch U. A note on the stability of a family of space-periodic Beltrami flows, *J. Fluid Mech.* **180**, 557–564 (1987).
- Glatzmaier G.A. and Roberts P.H. A three-dimensional self-consistent computer simulation of a geomagnetic field reversal. *Nature* **377**, 203–209 (1995).
- Glatzmaier G.A. and Roberts P.H. An anelastic geodynamo simulation driven by compositional and thermal convection, *Physica D* **97**, 81–94 (1996).

- Hide R., Skeldon A.C. and Acheson D.J. A study of two novel self-exiting single-disc homopolar dynamos: theory, *Proc. R. Soc. Lond. A* **452**, 1369–1395 (1996).
- Jacobs J.A. *Reversals of the Earth's Magnetic Field*, Cambridge University Press, 1994.
- Jones C.A. Convection driven geodynamo models, *Phil. Trans. R. Soc. Lond. A* **358**, 873–897 (2000).
- Matthews, P.C., Dynamo action in simple convective flows, *Proc. R. Soc.* **455**, 1829–1840 (1999).
- Melbourne, I., Proctor, M.R.E. and Rucklidge, A.M., A heteroclinic model of geodynamo reversals and excursions, In: *Dynamo and Dynamics, a Mathematical Challenge* (Eds. Chossat, P., Armbruster, D. and Oprea, I.), 363–370, Kluwer Academic Publishers (2001).
- Merrill R.T., McElhinny M.W. and Stevenson D.J. Evidence for long-term asymmetries in the Earth's magnetic field and possible implications for dynamo theories, *Phys. Earth Planet. Int.* **20**, 75–82 (1979).
- Merrill R.T. and McFadden P.L. Secular variation and the origin of geomagnetic field reversals, *J. Geophys. Res.* **93**, 11589–11597 (1986).
- Moroz I.M., Hide R. and Soward A.M. On self-exiting coupled Faraday disc homopolar dynamos driving series motors, *Physica D* **117**, 128–144 (1998).
- Podvigina O. and Pouquet A. On the non-linear stability of the 1:1:1 ABC flow, *Physica D* **75**, 471–508 (1994).
- Podvigina O.M. *Spatially-periodic evolutionary and steady solutions to the three-dimensional Navier-Stokes equation with the ABC-force*. Institute of Mechanics, Lomonosov Moscow State University (1999).
- Roberts P.H. and Glatzmaier G.A. The geodynamo , past, present and future, *Geophys. Astrophys. Fluid Dynamics* **94**, 47–84 (2001).
- Roberts P.H. and Stix M α -Effect Dynamos, by the Bullard-Gellman Formalism, *Astron. and Astrophys.* **18**, 453–466, (1972).
- Sarson G.R. and Jones C.A. A convection driven geodynamo reversal model, *Phys. Earth Planet. Int.* **111**, 3–20 (1999).
- Sarson G.R. Reversal models from dynamo calculations *Phil. Trans. Soc. Lond. A* **358**, 921–942 (2000).
- Zeldovich Ya.B., Ruzmaikin A.A. and Sokoloff D.D. *Magnetic fields in astrophysics*, Gordon and Breach, Science Publishers (1983).




Figure 1. Bifurcation diagram of the detected MHD attractors for $R = 4$ and $0 < R_m \leq 58$. For each type of attractors the interval of R_m 's (horizontal axis, non-uniform scale) is indicated, for which attractors of this type have been detected in computations. Labeling of attractors is explained in Section 4 (see also Table 1). Only one representative symmetry-related attractors is shown. The attractor $P_{2,1}^1$ for $R_m = 39.6$ and attractors from the second family are not shown. Stable steady states are represented by solid lines, the unstable ones – by dashed lines.




Figure 2a. Magnetic energy (vertical axis) as a function of time (horizontal axis) for $R = 4$ and $R_m = 30$.




Figure 2b. Projection of the trajectory in saturated regime (a periodic orbit $P_{1,i}$) on the plane of Fourier coefficients $\text{Im } b_{0,1,2}^1$ (horizontal axis) and $\text{Re } v_{0,1,2}^1$ (vertical axis) for $R = 4$ and $R_m = 30$ (same run as on Fig. 1a).




Figure 3a. Magnetic energy (vertical axis) as a function of time (horizontal axis) for $R = 4$ and $R_m = 39.2$.




Figure 3b. Projection of the trajectory in saturated regime (a periodic orbit $P_{2,i}^1$) on the plane of Fourier coefficients $\text{Im } b_{0,1,2}^1$ (horizontal axis) and $\text{Re } v_{0,1,2}^1$ (vertical axis) for $R = 4$ and $R_m = 39.2$ (same run as on Fig. 2a).




Figure 4a. Projection of the trajectory in saturated regime (a periodic orbit $P_{2,i}^2$) on the plane of Fourier coefficients $\text{Im } b_{0,1,2}^1$ (horizontal axis) and $\text{Re } v_{0,1,2}^1$ (vertical axis) for $R = 4$ and $R_m = 39.4$.




Figure 4b. Projection of the trajectory in saturated regime (a periodic orbit $P_{2,i}^4$) on the plane of Fourier coefficients $\text{Im } b_{0,1,2}^1$ (horizontal axis) and $\text{Re } v_{0,1,2}^1$ (vertical axis) for $R = 4$ and $R_m = 39.45$.




Figure 4c. Projection of the trajectory in the phase space in saturated regime (a periodic orbit $P_{2,i}^\infty$) on the plane of Fourier coefficients $\text{Im } b_{0,1,2}^1$ (horizontal axis) and $\text{Re } v_{0,1,2}^1$ (vertical axis) for $R = 4$ and $R_m = 39.5$.




Figure 5a. Projection of the trajectory in saturated regime (periodic orbit P_3) on the plane of Fourier coefficients $\text{Im } b_{0,1,2}^1$ (horizontal axis) and $\text{Re } v_{0,1,2}^1$ (vertical axis) for $R = 4$ and $R_m = 39.6$.




Figure 5b. Fourier coefficient $\text{Im } b_{0,1,2}^1$ (vertical axis) as a function of time (horizontal axis) for $R = 4$ and $R_m = 39.6$ (same run as on Fig. 4a).




Figure 5c. Magnetic energy (vertical axis) versus time (horizontal axis) for $R = 4$ and $R_m = 39.6$ (same run as on Fig. 4a).




Figure 5d. Kinetic energy (vertical axis) versus time (horizontal axis) for $R = 4$ and $R_m = 39.6$ (same run as on Fig. 4a).




Figure 6. Projection of the trajectory in saturated regime (periodic orbit P_4) on the plane of Fourier coefficients $\text{Im } b_{0,1,2}^1$ (horizontal axis) and $\text{Re } v_{0,1,2}^1$ (vertical axis) for $R = 4$ and $R_m = 39.7$.




Figure 7a. Fourier coefficient $\text{Im } b_{0,1,2}^1$ (vertical axis) as a function of time (horizontal axis) for $R = 4$ and $R_m = 42$.




Figure 7b. Fourier coefficient $\text{Re } b_{0,1,1}^3$ (vertical axis) as a function of time (horizontal axis) for $R = 4$ and $R_m = 42$ (same run as on Fig. 6a).




Figure 8a. Fourier coefficient $\text{Im } b_{0,1,2}^1$ (vertical axis) as a function of time (horizontal axis) for $R = 4$ and $R_m = 45$.




Figure 8b. Fourier coefficient $\text{Re } b_{0,1,1}^3$ (vertical axis) as a function of time (horizontal axis) for $R = 4$ and $R_m = 45$ (same run as on Fig. 7a).




Figure 9a. Fourier coefficient $\text{Im } b_{0,1,2}^1$ (vertical axis) as a function of time (horizontal axis) for $R = 4$ and $R_m = 46$.




Figure 9b. Fourier coefficient $\text{Re } b_{0,1,1}^3$ (vertical axis) as a function of time (horizontal axis) for $R = 4$ and $R_m = 46$ (same run as on Fig. 8a).




Figure 9c. Fourier coefficient $\text{Re } b_{1,0,1}^3$ (vertical axis) as a function of time (horizontal axis) for $R = 4$ and $R_m = 46$ (same run as on Fig. 8a).




Figure 10. Fourier coefficient $\text{Im } b_{0,1,2}^1$ (vertical axis) as a function of time (horizontal axis) for $R = 4$ and $R_m = 48$.




Figure 11a. Fourier coefficients $\text{Im } b_{0,1,2}^1$ (vertical axis) as a function of time (horizontal axis) for $R = 4$ and $R_m = 51$.




Figure 11b. Projection of the trajectory in saturated regime on the plane of Fourier coefficients $\text{Im } b_{0,1,2}^1$ (horizontal axis) and $\text{Re } v_{0,1,2}^1$ (vertical axis) for $R = 4$ and $R_m = 51$ (same run as on Fig. 10a).




Figure 12a. Fourier coefficient $\text{Im } b_{0,1,2}^1$ (vertical axis) as a function of time (horizontal axis) for $R = 4$ and $R_m = 52$.




Figure 12b. Projection of the trajectory in saturated regime (torus $T_{3,i}$) on the plane of Fourier coefficients $\text{Im } b_{0,1,2}^1$ (horizontal axis) and $\text{Re } v_{0,1,2}^1$ (vertical axis) for $R = 4$ and $R_m = 52$ (same run as on Fig. 11a).




Figure 13. Magnetic energy (vertical axis) as a function of time (horizontal axis) for $R = 15$ and $R_m = 40$.




Figure 14. Magnetic energy (vertical axis) as a function of time (horizontal axis) for $R = 20$ and $R_m = 40$.




Figure 15a. Kinetic energy (vertical axis) as a function of time (horizontal axis) for $R = 25$ and $R_m = 40$.




Figure 15b. Fourier coefficients of the flow (vertical axis: solid line – $\text{Re } v_{1,0,0}^3$, dashed line – $\text{Re } v_{0,1,0}^1$, dot line – $\text{Re } v_{0,0,1}^2$) as a function of time (horizontal axis) for $R = 25$ and $R_m = 40$ (same run as on Fig. 14a).




Figure 15c. Magnetic energy (vertical axis) as a function of time (horizontal axis) for $R = 25$ and $R_m = 40$ (same run as on Fig. 14a).




Figure 15d. Fourier coefficient $\text{Im } b_{0,1,2}^1$ (vertical axis) as a function of time (horizontal axis) for $R = 25$ and $R_m = 40$ (same run as on Fig. 14a).




Figure 16a. Kinetic energy (vertical axis) as a function of time (horizontal axis) for $R = 25$ in the hydrodynamic problem (the initial condition for the flow is the same as on Fig. 14).




Figure 16b. Fourier coefficients of the flow (vertical axis: solid line – $\text{Re } v_{1,0,0}^3$, dashed line – $\text{Re } v_{0,1,0}^1$, dot line – $\text{Re } v_{0,0,1}^2$) as a function of time (horizontal axis) for $R = 25$ (same run as on Fig. 15a).